

AVERAGED EXTREME REGRESSION QUANTILE

Jana Jurečková¹

¹ Charles University in Prague, Czech Republic

Consider the linear regression model

$$Y_{ni} = \mathbf{x}_{ni}^\top \boldsymbol{\beta} + e_{ni}, \quad i = 1, \dots, n$$

with observations Y_{n1}, \dots, Y_{nn} , independent errors e_{n1}, \dots, e_{nn} , identically distributed according to an unknown distribution function F ;

$\mathbf{x}_{ni} = (x_{i1}, \dots, x_{ip})^\top$ with $x_{i1} = 1$, $i = 1, \dots, n$ is the vector of covariates, and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)^\top$ is an unknown parameter. The *maximal regression quantile* $\widehat{\boldsymbol{\beta}}_n(1)$ is a solution of the minimization problem:

$$\min_{\mathbf{b} \in \mathbb{R}^p} \sum_{i=1}^n (Y_i - \mathbf{x}_i^\top \mathbf{b})^+,$$

which can be alternatively described as any solution to the linear program:

$$\min_{\mathbf{b} \in \mathbb{R}^p} \sum_{i=1}^n \mathbf{x}_i^\top \mathbf{b} \quad \text{s.t.} \quad Y_i \leq \mathbf{x}_i^\top \mathbf{b}, \quad i = 1, \dots, n.$$

We are interested in the averaged extreme regression quantile $\frac{1}{n} \sum_{i=1}^n \mathbf{x}_{ni}^\top \widehat{\boldsymbol{\beta}}_n(1)$ and in the relation of $\frac{1}{n} \sum_{i=1}^n \mathbf{x}_{ni}^\top (\widehat{\boldsymbol{\beta}}_n(1) - \boldsymbol{\beta})$ to the extreme quantiles of model errors e_{n1}, \dots, e_{nn} . Jurečková and Pícek (2014) studied the averaged regression α -quantile $\widehat{\boldsymbol{\beta}}_n(\alpha)$ for $0 < \alpha < 1$, and proved that

$$n^{1/2} [\bar{\mathbf{x}}_n^\top (\widehat{\boldsymbol{\beta}}_n(\alpha) - \boldsymbol{\beta}) - e_{n:[n\alpha]}] = \mathcal{O}_p(n^{-1/4}) \quad \text{as } n \rightarrow \infty.$$

The situation is different with the averaged extreme regression quantile. Rewrite the regression model in the form $\mathbf{Y}_n = \mathbf{X}_n \boldsymbol{\beta} + \mathbf{e}_n$, and consider the following linear programming problem, connected with $\widehat{\boldsymbol{\beta}}_n(1)$:

$$\begin{aligned} \mathbf{Y}_n^\top \mathbf{A} &= \max \\ \mathbf{X}_n^\top \mathbf{A} &= \mathbf{X}_n^\top \mathbf{1}_n \\ A_i &\geq 0, \quad i = 1, \dots, n \end{aligned}$$

with the optimal base \mathbf{X}_1 , a submatrix of \mathbf{X}_n , of order $p \times p$. Then

$$\bar{\mathbf{x}}_n^\top (\widehat{\boldsymbol{\beta}}_n(1) - \boldsymbol{\beta}) = \sum_{i=1}^p \lambda_i e_{n:n-i+1} \quad \text{for any finite } n > p,$$

where $e_{n:1} \leq \dots \leq e_{n:n}$ are order statistics corresponding to e_1, \dots, e_n and

$$\boldsymbol{\lambda}^\top = (\lambda_1, \dots, \lambda_p) = \frac{1}{n} \mathbf{1}_n^\top \mathbf{X}_n \mathbf{X}_1^{-1}.$$

The coefficients λ_i are obviously random and satisfy $\lambda_i > 0$, $i = 1, \dots, p$ and $\sum_{i=1}^p \lambda_i = 1$.

Keywords: α -regression quantile, Extreme regression quantile, Averaged extreme regression quantile

References:

- Jurečková, J. (2007). Remark on extreme regression quantile. *Sankhyā* 69, 87–100.
- Jurečková, J. and J. Picek (2005). Two-step regression quantiles. *Sankhya* 67, 227–252.
- Jurečková, J. and J. Picek (2014). Averaged regression quantiles. In: S. Lahiri et al., Eds.), *Contemporary Developments in Statistical Theory* (pp. 203–216). Springer Proc. in Math. & Statistics 68.
- Koenker, R. and G. Bassett (1978). Regression quantiles. *Econometrica* 46, 33–50.
- Portnoy, S. and J. Jurečková (1999). On extreme regression quantiles. *Extremes* 2:3, 227–243.
- Smith, R. (1994). Nonregular regression. *Biometrika* 81, 173–183.